

Shifted cotangent stacks are shifted symplectic

Damien Calaque

Abstract. We prove that shifted cotangent stacks carry a canonical shifted symplectic structure. We also prove that shifted conormal stacks carry a canonical Lagrangian structure. These results were believed to be true but no written proof was available in the Artin case.

Contents

1	Basics of derived (pre)symplectic geometry	2
1.1	Some ∞ -categorical linear algebra	2
1.2	Shifted (pre)symplectic, isotropic and Lagrangian structures	3
1.3	Isotropic and Lagrangian fibrations	4
2	The n-shifted symplectic structure of the n-shifted cotangent stack	5
2.1	An n -shifted presymplectic structure on $T^*[n]X$	6
2.2	Non-degeneracy of the n -shifted presymplectic structure	7
2.3	A Lagrangian structure on shifted conormal stacks	10
3	Shifted cotangent stacks in derived Poisson geometry	13
3.1	The n -shifted Poisson structure of $T^*[n]X$ and shifted polyvector fields	13
3.2	Lagrangian morphisms, deformed shifted cotangent stacks and shifted Poisson structures .	13

Introduction

Shifted symplectic and Lagrangian structures have been introduced in the inspiring paper [PTVV], where many examples of shifted symplectic stacks are given. They appeared to be very powerful tools, allowing for instance:

- to prove existence of symplectic structures on various moduli spaces ([PTVV]).
- to prove existence and functoriality of perfect symmetric obstruction theories ([PTVV]).
- to construct “classical” topological field theories ([Ca, Ha]).

Many results of classical symplectic geometry do extend to the shifted/derived context, sometimes even in a better way thanks to the flexibility of derived geometry. For instance, Lagrangian correspondences compose well, without any transversality assumption. But some obvious results happen to be much more difficult in the derived context, precisely because of the flexibility of derived geometry. The main example of this phenomenon is the classical (and easy) correspondence between symplectic and non-degenerate Poisson structures; in the shifted/derived setting this correspondence is much harder to prove, even in the affine case (see [CPT+, Pr]).

Another classical and rather easy fact in symplectic geometry is that the cotangent T^*Y to a manifold Y admits a canonical symplectic structure and that the conormal $N^*X \subset T^*Y$ of a submanifold $X \subset Y$ is Lagrangian. An analog of this was expected to be true for shifted cotangent (and conormal) stacks of derived Artin stacks, but no written proof was available (the Deligne-Mumford case is treated in [PTVV]).

We provide a proof in this paper. The spirit of the proof is essentially the same as the one in usual symplectic geometry. The challenge is to formulate it in a completely coordinate independent fashion.

We also conclude the paper with several conjectures about the relation between Lagrangian morphisms, (deformed) shifted cotangent stacks, and shifted Poisson structures in the sense of [CPT+, Pr].

Basic notation and conventions

- k is a field of characteristic zero (or just a noetherian \mathbb{Q} -algebra).
- all algebro-geometric structure are over k . For instance, a derived stack is a derived k -stack.
- a *cdga* means a commutative differential graded k -algebra sitting in non-positive degree. For commutative differential graded algebras without any degree condition we will say *unbounded cdga*.
- An Artin stack means a derived geometric k -stack locally of finite presentation in the sense of [HAG2].
- For a derived stack X and a quasi-coherent module E on X , we define $\mathbf{E}^*[n] := \mathbb{R}\mathrm{Spec}_X(\mathrm{Sym}(E[-n]))$ for every $n \in \mathbb{Z}$. We obviously write $\mathbf{E}^* = \mathbf{E}^*[0]$. Whenever E is perfect then $\mathbf{E}^*[n]$ is the derived stack of sections of the shifted dual $E^\vee[n]$ of E : an A -point of $\mathbf{E}^*[n]$, where A is a cdga, is an A -point $x : \mathbb{R}\mathrm{Spec}(A) \rightarrow X$ of X together with a section of $x^*E^\vee[n]$.
- In particular if X a derived Artin stack then its cotangent \mathbb{L}_X (relative to k) is perfect, and we denote by \mathbb{T}_X its dual. Hencefore we can define the n -shifted cotangent derived stack $\mathbf{T}^*[n]X$ of X as $\mathbf{T}^*[n]X := \mathbb{R}\mathrm{Spec}_X(\mathrm{Sym}(\mathbb{T}_X[-n]))$. We obviously write $\mathbf{T}^*X = \mathbf{T}^*[0]X$.
- similarly, we have $\mathbf{T}^*[n]f = \mathbb{R}\mathrm{Spec}_X(\mathrm{Sym}(\mathbb{T}_f[-n]))$ for a morphism $f : X \rightarrow Y$ of derived Artin stacks.
- as we work we ∞ -categories, unless otherwise specified all usual categorical terms must be understood ∞ -categorically (for instance, “limit” means “ ∞ -limit” or “homotopy limit”).

Acknowledgements

I thank Pavel Safronov for several stimulating discussions about shifted cotangent stacks. I actually first came up with a different (and quite indirect) strategy for the proof, that uses shifted symplectic groupoids (it will appear in a forthcoming work), and it was after a discussion with Pavel during the conference “Homotopical Methods in Quantum Field Theory” at IBS-CGP in Pohang that I started to look for a simpler and more direct proof.

I acknowledge the support of the Institut Universitaire de France and ANR SAT.

1 Basics of derived (pre)symplectic geometry

1.1 Some ∞ -categorical linear algebra

Let \mathcal{C} be a stable symmetric monoidal ∞ -category. Let V be a perfect (meaning dualizable) object in \mathcal{C} .

Definition 1.1. a) A degree n pairing $\omega : V^{\otimes 2} \rightarrow \mathbf{1}[n]$, where $\mathbf{1}$ is a unit, is *skew-symmetric* if it factors through $V^{\otimes 2} \rightarrow \wedge^2 V \rightarrow \mathbf{1}[n]$, where $\wedge^2 V := S^2(V[-1])[2]$. From now on, unless otherwise specified, all pairings will be skew-symmetric.

b) A degree n pairing $\omega : V^{\otimes 2} \rightarrow \mathbf{1}[n]$ is *non-degenerate* if the adjoint morphism $\omega^b : V \rightarrow V^\vee[n]$ is an equivalence.

Let $f : V \rightarrow W$ be a morphism between perfect objects in \mathcal{C} .

Definition 1.2. Let ω be a degree n pairing on V .

a) An *isotropic structure* on f (w.r.t. ω) is a homotopy between 0 and $f^*\omega$ in the space $\mathrm{Map}_{\mathcal{C}}(\wedge^2 U, \mathbf{1}[n])$ of degree n pairings on U .

b) An isotropic structure on f is *non-degenerate* if the induced null-homotopic sequence $U \rightarrow V \rightarrow U^\vee[n]$ is a fiber sequence.

The following result is very easy, though quite useful.

Lemma 1.3. *Assume that we are given a morphism $f : \mathbf{V} \rightarrow \mathbf{W}$ between perfect objects together with an n -shifted pairing ω on \mathbf{V} and an isotropic structure γ on f . If γ is non-degenerate then ω is non-degenerate as well.*

Proof. By definition, the null-homotopic sequence $\mathbf{U} \rightarrow \mathbf{V} \rightarrow \mathbf{U}^\vee[n]$ is a fiber sequence. Its (shifted) dual $\mathbf{U} \rightarrow \mathbf{V}^\vee[n] \rightarrow \mathbf{U}^\vee[n]$ is thus a fiber sequence as well. The main point is that these two sequences are mapped to each other in the following way:

$$\begin{array}{ccccc} \mathbf{U} & \longrightarrow & \mathbf{V} & \longrightarrow & \mathbf{V}^\vee[n] \\ \parallel & & \downarrow \omega^\flat & & \parallel \\ \mathbf{U} & \longrightarrow & \mathbf{V}^\vee[n] & \longrightarrow & \mathbf{V}^\vee[n] \end{array}$$

Hence the morphism $\omega^\flat : \mathbf{E} \rightarrow \mathbf{E}^\vee[n]$ is an equivalence and ω is non-degenerate. \square

1.2 Shifted (pre)symplectic, isotropic and Lagrangian structures

Recall from [PTVV] that to any derived stack X one can associate a graded mixte module $\mathbf{DR}(X)$. We will often denote d_{dR} the mixte differential ϵ (of weight 1 and cohomological degree 1)¹, and d_{int} or d_X the differential of the underlying graded module $\mathbf{DR}(X)^\sharp$ (of weight 0 and cohomological degree 1), also called the *internal differential*.

If $X = \text{Spec}(A)$, for a cofibrant cdga A , then $\mathbf{DR}(X) = \mathbf{DR}(A/\mathbf{k})$ is the graded module

$$\text{Sym}_A(\Omega_A^1-1).$$

equipped with the following mixte differential:

$$\epsilon = d_{\text{dR}} : a_0(da_1) \dots (da_n) \mapsto (da_0)(da_1) \dots (da_n).$$

Note that (da_i) is *a priori* not “ d applied to a_i and is in particular **not** $d_{\text{int}}a_i = d_Aa_i$. But observe that $(da_i) = d_{\text{dR}}a_i$; hence we will try to use the notation $d_{\text{dR}}a_i$ in order to avoid confusion.

Definition 1.4. a) The space of *p-forms of degree n on X* is

$$A^p(X, n) := \text{Map}_{\mathbf{k}\text{-mod}^{\text{gr}}}(\mathbf{k}, \mathbf{DR}(X)^\sharp[n+p](p)).$$

b) The space of *closed p-forms of degree n on X* is

$$A^{p, \text{cl}}(X, n) := \text{Map}_{\mathbf{k}\text{-mod}^{\text{gr}, \text{mixt}}}(\mathbf{k}, \mathbf{DR}(X)[n+p](p)).$$

c) For a closed p -form of α degree n , we denote by α_0 its *underlying p-form of degree n* , that is to say its image through the forgetful functor $(-)^{\sharp}$.

The functors $A^p(n) := A^p(-, n)$ and $A^{p, \text{cl}}(n) := A^{p, \text{cl}}(-, n)$ are actually derived stacks (see [PTVV, Proposition 1.11]).

Remark 1.1. Note that for a given graded mixte module M , $\text{Map}_{\mathbf{k}\text{-mod}^{\text{gr}, \text{mixt}}}(\mathbf{k}, M)$ is the space of a semi-infinite sequences

$$\alpha = (\alpha_0, \dots, \alpha_i, \dots)$$

such that $d(\alpha_0) = 0$, and for every $i \geq 0$, $\epsilon(\alpha_i) = d(\alpha_{i+1})$. I.e. it is the space of 0-cocycles in $\prod_{k \geq 0} M^{(p)}$ equipped with the total differential $d + \epsilon$. The image of such a sequence α through the forgetful functor $(-)^{\sharp}$ is precisely its leading term α_0 , which is a 0-cocyle in $M^{(0)}$. We will thus sometimes write, in the context of the above definition, “*the leading term*” in place of “the underlying p -form of degree n ” of a closed p -form of degree n .

¹We use the weight/grading convention from [CPT+] rather than the one of [PTVV].

Definition 1.5. a) An \mathfrak{n} -shifted presymplectic structure on a stack X is a closed 2-form of degree \mathfrak{n} on X .

b) Let (X, ω) be a stack equipped with an \mathfrak{n} -shifted presymplectic structure. An *isotropic structure* on a morphism $f : L \rightarrow X$ is a path from 0 to $f^*\omega$ in the space $A^{2,cl}(L, \mathfrak{n})$ of \mathfrak{n} -shifted presymplectic structures on L .

c) We say that an \mathfrak{n} -shifted presymplectic structure ω on an X is *symplectic* if X has a perfect cotangent complex \mathbb{L}_X and if the degree \mathfrak{n} pairing on $\mathbb{T}_X := \mathbb{L}_X^\vee$ given by its leading term ω_0 is non-degenerate in the sense of Definition 1.1.

d) We say that an isotropic structure γ on $f : L \rightarrow X$ is *non-degenerate* if both X and L have perfect cotangent complexes and if the leading term γ_0 is indeed a non-degenerate isotropic structure on the morphism $\mathbb{T}_L \rightarrow f^*\mathbb{T}_X$ in the sense of Definition 1.2. It is only when the presymplectic structure ω itself is symplectic that we name *Lagrangian structures* the non-degenerate isotropic ones.

The main result of Section 2 below is that the \mathfrak{n} -shifted cotangent stack $\mathbf{T}^*[\mathfrak{n}]X$ of an Artin stack X carries an \mathfrak{n} -shifted symplectic structure and that the zero section morphism $X \rightarrow \mathbf{T}^*[\mathfrak{n}]X$ carries a Lagrangian structure.

We refer to [PTVV, Ca, Sa] for many other examples of \mathfrak{n} -shifted symplectic and Lagrangian structures.

Remark 1.2. a) A stack equipped an \mathfrak{n} -shifted presymplectic structure (an \mathfrak{n} -shifted presymplectic stack, for short) is nothing but a stack over $A^{2,cl}(\mathfrak{n})$.

b) Let (X, ω) be a stack equipped with an \mathfrak{n} -shifted presymplectic structure. An *isotropic structure* on a morphism $f : L \rightarrow X$ is nothing but the data of a commuting square

$$\begin{array}{ccc} L & \xrightarrow{f} & X \\ \downarrow & & \downarrow \omega \\ * & \xrightarrow{0} & A^{2,cl}(\mathfrak{n}) \end{array}$$

The above alternative approach to isotropic structures on morphisms (due to Rune Haugseng [Ha, Section 9]) leads to the following:

Definition 1.6. An \mathfrak{n} -isotropic correspondence is a correspondence in the ∞ -category $\mathbf{dSt}_{/A^{2,cl}(\mathfrak{n})}$ of derived stacks over $A^{2,cl}(\mathfrak{n})$: it is the data of a commuting square

$$\begin{array}{ccc} L & \xrightarrow{g} & Y \\ \downarrow f & & \downarrow \omega_Y \\ X & \xrightarrow{\omega_X} & A^{2,cl}(\mathfrak{n}) \end{array}$$

In other words, an \mathfrak{n} -isotropic correspondence is the data of two \mathfrak{n} -shifted presymplectic stacks (X, ω_X) and (Y, ω_Y) together with an isotropic morphism $f \times g : L \rightarrow \overline{X} \times Y$, where $\overline{X} \times Y$ is the \mathfrak{n} -shifted presymplectic stack $(X \times Y, \pi_2^* \omega_Y - \pi_1^* \omega_X)$. Assuming ω_X and ω_Y are symplectic, then $f \times g$ is Lagrangian if and only if the commuting square

$$\begin{array}{ccc} \mathbb{T}_L & \longrightarrow & g^* \mathbb{T}_Y \simeq g^* \mathbb{L}_Y[\mathfrak{n}] \\ \downarrow & & \downarrow \\ g^* \mathbb{T}_X \simeq f^* \mathbb{L}_X[\mathfrak{n}] & \longrightarrow & \mathbb{L}_L[\mathfrak{n}] \end{array}$$

is (co)Cartesian. In this case we talk about an \mathfrak{n} -Lagrangian correspondence.

1.3 Isotropic and Lagrangian fibrations

Recall from [CPT+, Section 1] that to any morphism $f : X \rightarrow Y$ of derived stacks one can associate a graded mixte module $\mathbf{DR}(f) := \mathbf{DR}(X/Y)$. If $f : X = \mathrm{Spec}(A) \rightarrow \mathrm{Spec}(B) = Y$ comes from a cofibration

$B \rightarrow A$ between cofibrant cdgas, then $\mathbf{DR}(X/Y) := \mathbf{DR}(A/B)$ is the graded module

$$\mathrm{Sym}_A(\Omega_{A/B}^1-1).$$

equipped with the mixte differential:

$$\epsilon : a_0(da_1) \dots (da_n) \mapsto (da_0)(da_1) \dots (da_n).$$

Definition 1.7. a) The space of *relative p-forms of degree n on f* (a-k-a *p-forms of degree n on X relative to Y*) is

$$A^p(f, n) = A^p(X/Y, n) := \mathrm{Map}_{\mathbf{k}\text{-mod}^{gr}}(\mathbf{k}, \mathbf{DR}(X/Y)^\sharp[n+p](p)).$$

b) The space of *relative closed p-forms of degree n on f* is

$$A^{p, cl}(f, n) = A^{p, cl}(X/Y, n) := \mathrm{Map}_{\mathbf{k}\text{-mod}^{gr, mixt}}(\mathbf{k}, \mathbf{DR}(X/Y)[n+p](p)).$$

c) For a relative closed p-form of α degree n, we denote by α_0 its *underlying relative p-form of degree n*, that is to say its image through the forgetful functor $(-)^{\sharp}$.

Note that we have a null-homotopic sequence

$$\mathbf{DR}(Y) \xrightarrow{f^*} \mathbf{DR}(X) \xrightarrow{-/Y} \mathbf{DR}(X/Y).$$

Remark 1.3. Recall from [CPT+, Remark 2.4.8] that $\mathbf{DR}(X/Y)$ can be obtained as global sections of a stack $\mathcal{DR}(f) = \mathcal{DR}(X/Y)$ of graded mixte modules over the ∞ -site of derived affine schemes over Y :

$$\mathcal{DR}(X/Y)(\mathrm{Spec}(A) \xrightarrow{y} Y) := \mathbf{DR}(y^*X/\mathrm{Spec}(A)).$$

Observe that $\mathcal{DR}(X/Y)$ is actually a graded mixed \mathcal{O}_Y -module as every $\mathbf{DR}(y^*X/\mathrm{Spec}(A))$ is a graded mixte A -module.

Let now X be equipped with an n -shifted presymplectic structure ω .

Definition 1.8. a) An *isotropic fibration structure on f* is a path from 0 to ω/Y in the space $A^{2, cl}(X/Y, n)$ of relative closed 2-forms of degree n .

b) An isotropic fibration structure γ on f is *non-degenerate* (or is a *Lagrangian fibration structure*) if both X and Y have perfect cotangent complexes and if the leading term γ_0 is indeed a non-degenerate isotropic structure on the morphism $\mathbb{T}_{X/Y} \rightarrow \mathbb{T}_X$ in the sense of Definition 1.2.

Remark 1.4. It follows from Lemma 1.3 that if γ is non-degenerate then the n -shifted presymplectic structure ω is indeed n -shifted symplectic. Therefore we don't need here to make a distinction between non-degenerate isotropic fibration structures and Lagrangian fibration structures.

We will see in Section 2 that the projection $\mathbf{T}^*[n]X \rightarrow X$ of a shifted cotangent stack (of an Artin stack) comes equipped with a Lagrangian fibration structure.

2 The n -shifted symplectic structure of the n -shifted cotangent stack

Let X be an Artin stack. It has a perfect global *cotangent complex* \mathbb{L}_X (see [HAG2]), its dual $\mathbb{T}_X := \mathbb{L}_X^\vee$ being called the *tangent complex*. Our main object of interest is its n -shifted cotangent stack $\mathbf{T}^*[n]X := \mathbb{R}\mathrm{Spec}_X(\mathrm{Sym}(\mathbb{T}_X[-n]))$, that is nothing but the stack of n -shifted 1-forms on X . Note that, given a perfect module E over X , the stack of sections $\mathbf{E} := A_X(E) := \mathbb{R}\mathrm{Spec}_X(\mathrm{Sym}(E^\vee))$ is an Artin stack as well (it is an affine stack of finite presentation over X).

In this Section we prove that the n -shifted cotangent stack $\mathbf{T}^*[n]X$ is n -shifted symplectic, together with variations on this result. All this was mainly already known for Deligne-Mumford stacks.

2.1 An n -shifted presymplectic structure on $T^*[n]X$

Recall that the space of morphisms $Y \rightarrow T^*[n]X$ is nothing but the space of morphisms $f : Y \rightarrow X$ equipped with a section of $f^*\mathbb{L}_X[n]$. In particular the identity map $T^*[n]X \rightarrow T^*[n]X$ corresponds to the data of the projection map $\pi_X : T^*[n]X \rightarrow X$ together with a section of $\pi_X^*\mathbb{L}_X[n]$. Therefore, using the map $\pi_X^*\mathbb{L}_X[n] \rightarrow \mathbb{L}_{T^*[n]X}[n]$ we obtain that $T^*[n]X$ carries a tautological n -shifted 1-form λ_X . The morphism $d_{\text{dR}} : A^1(-, n) \rightarrow A^{2, c^1}(-, n)$ sends it to an n -shifted closed 2-form $\omega_X = d_{\text{dR}}\lambda_X$ on $T^*[n]X$.

The zero section morphism $\iota_X : X \rightarrow T^*[n]X$ corresponds to the data of the identity map $X \rightarrow X$ together with the zero section of $\mathbb{L}_X[n]$. Tautologically, this means that the pull-back of λ_X along the zero section is homotopic to zero. Hence we obtain an isotropic structure γ_X on ι_X (w.r.t. the n -shifted presymplectic structure previously described).

Finally observe λ_X comes from a section of $\pi_X^*\mathbb{L}_X[n]$ and recall that the sequence

$$\pi_X^*\mathbb{L}_X \rightarrow \mathbb{L}_{T^*[n]X} \rightarrow \mathbb{L}_{\pi_X}$$

is fibered. Therefore the image of λ_X into relative n -shifted 1-forms along π_X is homotopic to zero. Hence we obtain that the projection morphism π_X is equipped with an isotropic fibration structure η_X .

Remark 2.1. Note that \mathbb{L}_{π_X} is nothing but $\pi_X^*\mathbb{T}_X[-n]$.

2.1.1 Compatibility with the \mathbb{G}_m -action

Observe that, for a perfect module E over X , its stack of sections \mathbf{E} is acted on by \mathbb{G}_m because $\text{Sym}(E^\vee)$ is a graded \mathcal{O}_X -algebra (by symmetric powers). We will refer to this new grading as the *fiber grading* (not to confuse it with the weight of forms and closed forms). Both the zero section $X \rightarrow \mathbf{E}$ and the projection $\mathbf{E} \rightarrow X$ are \mathbb{G}_m -equivariant for the trivial \mathbb{G}_m -action on X .

Notation 2.1. a) If Y is an Artin stack equipped with an action of \mathbb{G}_m , then the relative cotangent complex $\mathbb{L}_{[Y/\mathbb{G}_m]/B\mathbb{G}_m}$ is a quasi-coherent sheaf on $[Y/\mathbb{G}_m]$. Its pull-back along the quotient map $Y \rightarrow [Y/\mathbb{G}_m]$ identifies with \mathbb{L}_Y . We will therefore allow ourselves to abuse notation and to keep writing \mathbb{L}_Y for its \mathbb{G}_m -equivariant enhancement $\mathbb{L}_{[Y/\mathbb{G}_m]/B\mathbb{G}_m}$. Similarly, if $f : Y_1 \rightarrow Y_2$ is a \mathbb{G}_m -equivariant map, then we will keep writing \mathbb{L}_f for its \mathbb{G}_m -equivariant enhancement $\mathbb{L}_{[f]}$, where $[f] : [Y_1/\mathbb{G}_m] \rightarrow [Y_2/\mathbb{G}_m]$ is the induced map on quotient stacks.

b) Sheaves on $B\mathbb{G}_m = [*/\mathbb{G}_m]$ are identified with graded complexes. For a graded complex $M = (M_i)_{i \in \mathbb{Z}}$ we denote by $M\{k\}$ its k -th shift for this grading (meaning that $M\{k\}_i = M_{i+k}$). Without further precision, a genuine complex is always assumed to be equipped with the trivial \mathbb{G}_m -action. This notation extends as well to sheaves on $[X/\mathbb{G}_m] = X \times B\mathbb{G}_m$, that are nothing but graded sheaves on X .

Remark 2.2. With this notation the identification from Remark 2.1 becomes $\mathbb{L}_{\pi_X} \cong \pi_X^*\mathbb{T}_X[-n]\{-1\}$.

Now, if Y is a derived stack equipped with an action of \mathbb{G}_m , observe that a \mathbb{G}_m -equivariant map $Y \rightarrow \mathbf{E}$ is determined by the data of a \mathbb{G}_m -equivariant map $f : Y \rightarrow X$ together with a \mathbb{G}_m -equivariant section of $f^*\mathbf{E}\{1\}$. Applying this to the identity map $T^*[n]X \rightarrow T^*[n]X$ we get a \mathbb{G}_m -equivariant section of $\pi_X^*\mathbb{L}_X\{1\}$. As a consequence, the 1-form $\lambda_X \in A^1(T^*[n]X, n) \cong \Gamma(T^*[n]X, \mathbb{L}_{T^*[n]X}[n])$ of (cohomological) degree n is homogeneous of weight 1 for the fiber grading.

Recall from Remark 1.3 that for a stack Y equipped with a \mathbb{G}_m -action we have a sheaf $\mathcal{DR}([Y/\mathbb{G}_m]/B\mathbb{G}_m)$ of graded mixed \mathcal{O} -modules on $B\mathbb{G}_m$. Its pull-back along the canonical point in $B\mathbb{G}_m$ is $\mathbf{DR}(Y)$. This means that $\mathbf{DR}(Y)$ carries yet another auxiliary grading that enhances it to a graded mixed graded k -module. We will follow the (abusing) Notation 2.1 and keep writing $\mathbf{DR}(Y)$ for its graded enhancement $\mathcal{DR}([Y/\mathbb{G}_m]/B\mathbb{G}_m)$. Similarly, if $f : Y_1 \rightarrow Y_2$ is a \mathbb{G}_m -equivariant map, then $\mathbf{DR}(f)$ also admits \mathbb{G}_m -equivariant enhancement that we still denote the same.

Going back to $T^*[n]X$ we get that λ_X belongs to the space

$$A^{1, \{1\}}(T^*[n]X, n) := \text{Map}_{k\text{-mod}^{gr, gr}}(k, \mathbf{DR}(T^*[n]X)^\sharp[n+1](1)\{1\})$$

and that $\omega_X = d_{\text{dR}}\lambda_X$ belongs to the space

$$A^{2, \text{cl}, \{1\}}(\mathbf{T}^*[n]X, \mathfrak{n}) := \text{Map}_{\mathbf{k}\text{-mod}^{\text{gr}, \text{gr}, \text{mult}}}(\mathbf{k}, \mathbf{DR}(\mathbf{T}^*[n]X)[n+2](2)\{1\})$$

of closed 2-forms of (cohomological) degree \mathfrak{n} on $\mathbf{T}^*[n]X$ that are homogeneous of weight 1 for the fiber grading. Similarly, η_X is a path in the space

$$A^{2, \text{cl}, \{1\}}(\pi_X, \mathfrak{n}) := \text{Map}_{\mathbf{k}\text{-mod}^{\text{gr}, \text{gr}, \text{mult}}}(\mathbf{k}, \mathbf{DR}(\pi_X)[n+2](2)\{1\})$$

of relative closed 2-forms of (cohomological) degree \mathfrak{n} along π_X that are homogeneous of weight 1 for the fiber grading.

2.2 Non-degeneracy of the \mathfrak{n} -shifted presymplectic structure

We will now prove that ω_X actually defines an \mathfrak{n} -shifted symplectic structure on $\mathbf{T}^*[n]X$, that is to say the morphism

$$\mathbb{T}_{\mathbf{T}^*[n]X} \longrightarrow \mathbb{L}_{\mathbf{T}^*[n]X}[n]$$

induced by its underlying \mathfrak{n} -shifted 2-form is an equivalence.

We also prove that the isotropic morphism $X \rightarrow \mathbf{T}^*[n]X$ and the isotropic fibration $\mathbf{T}^*[n]X \rightarrow X$ are Lagrangian.

Theorem 2.2. *The \mathfrak{n} -shifted cotangent stack $\mathbf{T}^*[n]X$ is \mathfrak{n} -shifted symplectic, its zero section is Lagrangian, and the projection down to X is a Lagrangian fibration. More precisely:*

1. *The \mathfrak{n} -shifted presymplectic structure ω_X on $\mathbf{T}^*[n]X$ is non-degenerate.*
2. *The \mathfrak{n} -shifted isotropic structure γ_X on the zero section $\iota_X : X \rightarrow \mathbf{T}^*[n]X$ is non-degenerate.*
3. *The \mathfrak{n} -shifted isotropic fibration structure η_X on $\pi_X : \mathbf{T}^*[n] \rightarrow X$ is non-degenerate.*

Before we start it is important to have in mind the two following fiber sequences, that are (\mathfrak{n} -shifted) dual to each other:

$$\pi_X^* \mathbb{L}_X[n] \cong \mathbb{T}_{\pi_X} \rightarrow \mathbb{T}_{\mathbf{T}^*[n]X} \rightarrow \pi_X^* \mathbb{T}_X \quad \text{and} \quad \pi_X^* \mathbb{L}_X[n] \rightarrow \mathbb{L}_{\mathbf{T}^*[n]X}[n] \rightarrow \mathbb{L}_{\pi_X}[n] \cong \pi_X^* \mathbb{T}_X.$$

Moreover, when pulled-back along the zero section these sequences split and we get that

$$\iota_X^* \mathbb{T}_{\mathbf{T}^*[n]X} \cong \mathbb{L}_X[n] \oplus \mathbb{T}_X \quad \text{and} \quad \iota_X^* \mathbb{L}_{\mathbf{T}^*[n]X}[n] \cong \mathbb{L}_X[n] \oplus \mathbb{T}_X.$$

Remark 2.3. The above fiber sequences actually are \mathbb{G}_m -equivariant. More precisely, according to Remark 2.2 we thus get fiber sequences

$$\pi_X^* \mathbb{L}_X[n]\{1\} \rightarrow \mathbb{T}_{\mathbf{T}^*[n]X} \rightarrow \pi_X^* \mathbb{T}_X \quad \text{and} \quad \pi_X^* \mathbb{L}_X[n] \rightarrow \mathbb{L}_{\mathbf{T}^*[n]X}[n] \rightarrow \mathbb{L}_{\pi_X}[n] \cong \pi_X^* \mathbb{T}_X\{-1\}$$

of \mathbb{G}_m -equivariant sheaves on $\mathbf{T}^*[n]X$ that split when pulled-back along the zero section:

$$\iota_X^* \mathbb{T}_{\mathbf{T}^*[n]X} \cong \mathbb{L}_X[n]\{1\} \oplus \mathbb{T}_X \quad \text{and} \quad \iota_X^* \mathbb{L}_{\mathbf{T}^*[n]X}[n] \cong \mathbb{L}_X[n] \oplus \mathbb{T}_X\{-1\}.$$

2.2.1 The isotropic structure on the zero section ι_X is non-degenerate

We prove here the second part of Theorem 2.2 .

We have an isotropic structure γ_X on the zero section morphism $\iota_X : X \rightarrow \mathbf{T}^*[n]X$, inducing in particular a commuting diagram

$$\begin{array}{ccccc} \mathbb{T}_X & \xrightarrow{\quad} & \mathbb{T}_X & \xrightarrow{\quad} & 0 \\ \downarrow & & \downarrow & & \downarrow \\ \iota_X^* \mathbb{T}_{\mathbf{T}^*[n]X} & \longrightarrow & \iota_X^* \mathbb{L}_{\mathbf{T}^*[n]X}[n] & \longrightarrow & \mathbb{L}_X[n] \end{array}$$

in which the right-hand square is cartesian.

We will now show that

Lemma 2.3. *The morphism $\phi_X : \mathbb{T}_X \rightarrow \mathbb{T}_X$ appearing in the above diagram is natural in X , in the sense that for every morphism $f : X \rightarrow Y$ we get a commuting square*

$$\begin{array}{ccc} \mathbb{T}_X & \xrightarrow{\phi_X} & \mathbb{T}_X \\ \downarrow & & \downarrow \\ f^*\mathbb{T}_Y & \xrightarrow{f^*\phi_Y} & f^*\mathbb{T}_Y \end{array}$$

Corollary 2.4. *The isotropic structure on the zero section $\iota_Y : Y \rightarrow \mathbb{T}^*[n]Y$ is non-degenerate, that is to say the map ϕ_Y is an equivalence.*

Proof of the Corollary. We know that it is true for derived affine schemes locally of finite presentation. Let us then assume by induction that we know the result holds for m -Artin stacks (i.e. Artin stacks that are m -geometric in the sense of [HAG2]). Let us then prove that it is true as well for $(m+1)$ -Artin stacks. If Y is an $(m+1)$ -Artin stack then $Y = |G_*| := \operatorname{colim}_n (G_n)$, where $G_* = (G_n)_{n \geq 0}$ is a smooth Segal groupoid in m -Artin stacks. Let $q : X = G_0 \rightarrow Y$ be the natural map to the colimit. Note that $q^*\mathbb{T}_Y$ can be computed as the colimit of the simplicial diagram $(e_n^*\mathbb{T}_{G_n})_{n \geq 0}$, where $e_n : G_0 \rightarrow G_n$ is the unit map.

Thanks to the above Lemma we have a simplicial diagram $(e_n^*\phi_{G_n})_{n \geq 0}$ of morphisms of perfect modules on X , the colimit of which is the morphism $q^*\phi_Y$. Thanks to the induction hypothesis every $e_n^*\phi_{G_n}$ is an equivalence, and thus $q^*\phi_Y$ is an equivalence as well. One finally gets that ϕ_Y is an equivalence because q^* is conservative (q is a groupoid quotient morphism).

We get the result by induction on the degree of geometricity. \square

Remark 2.4. One can even prove that the equivalence ϕ_X is homotopic to the identity (this is true for derived scheme, and then the inductive proof is exactly the same).

Proof of the Lemma. Let $f : X \rightarrow Y$ be a map of Artin stacks and consider the diagram

$$\begin{array}{ccccc} & & X & \xrightarrow{f} & Y \\ & \swarrow & \downarrow & & \downarrow \iota_Y \\ \mathbb{T}^*[n]X & \longleftarrow & f^*\mathbb{T}^*[n]Y & \longrightarrow & \mathbb{T}^*[n]Y \end{array}$$

where $f^*\mathbb{T}^*[n]Y := \mathbb{T}^*[n]Y \times_Y X$ is the stack of sections of $f^*\mathbb{L}_Y[n]$.

The morphism ϕ_X , resp. ϕ_Y , is determined by the leading term of the isotropic structure γ_X , resp. γ_Y . Hence the morphism $f^*\phi_Y$ is determined by the leading term of $f^*\gamma_Y$, considered as an isotropic structure on

$$f^*\iota_Y : X \longrightarrow f^*\mathbb{T}^*[n]Y$$

and where we equip $f^*\mathbb{T}^*[n]Y$ with the n -shifted presymplectic structure $\tilde{\omega}$ given by the pull-back of ω_Y along $f^*\mathbb{T}^*[n]Y \rightarrow \mathbb{T}^*[n]Y$. Now the result follows from the following fact:

- the pull-back of the pair (ω_X, γ_X) along $f^*\mathbb{T}^*[n]Y \rightarrow \mathbb{T}^*[n]X$ is naturally homotopic to the pair $(\tilde{\omega}, f^*\gamma_Y)$.

Actually, the pull-back of λ_X itself along $f^*\mathbb{T}^*[n]Y \rightarrow \mathbb{T}^*[n]X$ is naturally homotopic to the pull-back of λ_Y along $f^*\mathbb{T}^*[n]Y \rightarrow \mathbb{T}^*[n]Y^2$. All diagrams we have written down live under X , therefore all previously considered homotopies between 0 and pull-backs of λ 's on X are themselves equivalent. The Lemma is proved. \square

²This is a slight variation on the obvious fact that for any morphism $u : A \rightarrow B$, $\operatorname{id}_B \circ u$ and $u \circ \operatorname{id}_A$ are equivalent.

2.2.2 The isotropic fibration structure on the projection π_X is non-degenerate

We now give the proof of the third part of the theorem. Its structure is very similar to the one of the second part (apart from that we will have to use \mathbb{G}_m -equivariant enhancements at some point).

We have an isotropic fibration structure η_X on the projection $\pi_X : \mathbf{T}^*[n]X \rightarrow X$, inducing in particular a commuting diagram

$$\begin{array}{ccccc} \pi_X^* \mathbb{L}_X[n] & \longrightarrow & \pi_X^* \mathbb{L}_X[n] & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{T}_{\mathbf{T}^*[n]X} & \longrightarrow & \mathbb{L}_{\mathbf{T}^*[n]X}[n] & \longrightarrow & \pi_X^* \mathbb{T}_X \end{array}$$

in which the right hand square is cartesian.

Let us write $\psi_X : \pi_X^* \mathbb{L}_X \rightarrow \pi_X^* \mathbb{L}_X$ for the $(-n)$ -shift of the morphism $\pi_X^* \mathbb{L}_X[n] \rightarrow \pi_X^* \mathbb{L}_X[n]$ appearing in the above diagram. Our aim is to prove that ψ_X is an equivalence. It follows from the next Lemma that it is sufficient to prove that $\iota_X^* \psi_X : \mathbb{L}_X \rightarrow \mathbb{L}_X$ is an equivalence.

Lemma 2.5. $\psi_X = \pi_X^* \iota_X^* \psi_X$.

Proof. Recall that, up to a shift, the morphism ψ_X is the composition of the map $\mathbb{T}_{\pi_X} \rightarrow \pi_X^* \mathbb{L}_X[n]$ given by the (leading term of the) isotropic fibration structure η_X with the canonical identification $\pi_X^* \mathbb{L}_X[n] \cong \mathbb{T}_{\pi_X}$ (dual to the one from Remark 2.1). Taking into account \mathbb{G}_m -equivariant enhancements, it becomes the composition of:

- the map $\mathbb{T}_{\pi_X} \rightarrow \pi_X^* \mathbb{L}_X[n]\{1\}$ given by (the leading term of the) path η that is homogeneous of weight 1 for the fiber grading,
- with the identification $\pi_X^* \mathbb{L}_X[n]\{1\} \cong \mathbb{T}_{\pi_X}$ (dual to the one from Remark 2.2).

Hence ψ_X is a \mathbb{G}_m -equivariant map $\pi_X^* \mathbb{L}_X\{1\} \rightarrow \pi_X^* \mathbb{L}_X\{1\}$. Since $\mathbb{L}_X\{1\}$ is a graded sheaf of pure weight on X , then ψ_X is the pull-back along π_X of a morphism $\mathbb{L}_X \rightarrow \mathbb{L}_X$ (that has to be $\iota_X^* \psi_X$). \square

Let now $f : X \rightarrow Y$ be a map of Artin stacks and consider the diagram

$$\begin{array}{ccccc} \mathbf{T}^*[n]X & \xleftarrow{\mathbf{T}^*f} & f^* \mathbf{T}^*[n]Y & \xrightarrow{\pi_Y^* f} & \mathbf{T}^*[n]Y \\ & \searrow \pi_X & \downarrow f^* \pi_Y & & \downarrow \pi_Y \\ & & X & \xrightarrow{f} & Y \end{array}$$

We will show that

Lemma 2.6. *The morphism $\psi_X : \pi_X^* \mathbb{L}_X \rightarrow \pi_X^* \mathbb{L}_X$ is natural in X , in the sense that for every morphism $f : X \rightarrow Y$ we get a commuting square*

$$\begin{array}{ccc} (f^* \pi_Y)^* \mathbb{L}_X & \xrightarrow{(\mathbf{T}^* f)^* \psi_X} & (f^* \pi_Y)^* \mathbb{L}_X \\ \uparrow & & \uparrow \\ (f^* \pi_Y)^* f^* \mathbb{L}_Y & \xrightarrow{(\pi_Y^* f)^* \psi_Y} & (f^* \pi_Y)^* f^* \mathbb{L}_Y \end{array}$$

In particular we have a commuting square

$$\begin{array}{ccc} \mathbb{L}_X & \xrightarrow{\iota_X^* \psi_X} & \mathbb{L}_X \\ \uparrow & & \uparrow \\ f^* \mathbb{L}_Y & \xrightarrow{f^* \iota_Y^* \psi_Y} & f^* \mathbb{L}_Y \end{array}$$

Corollary 2.7. *The isotropic fibration structure on the projection $\pi_X : T^*[n]X \rightarrow X$ is non-degenerate, that is to say the map ψ_X is an equivalence.*

Proof of the Corollary. It follows from Lemma 2.5 that it is sufficient to show that $\iota_X^* \psi_X$ is an equivalence, the proof of which is almost identical to the one of Corollary 2.4. Let us repeat it for the sake of clarity.

We know that it is true for derived affine schemes locally of finite presentation. Let us then assume by induction that we know the result holds for m -Artin stacks and prove that it does for $(m+1)$ -Artin stacks as well. If Y is an $(m+1)$ -Artin stack then $Y = |G_*|$, where G_* is a smooth Segal groupoid in m -Artin stacks. Let $q : X = G_0 \rightarrow Y$ be the natural map to the colimit. Note that $q^* \mathbb{L}_Y$ can be computed as the limit of the cosimplicial diagram $(e_n^* \mathbb{L}_{G_n})_{n \geq 0}$, where $e_n : G_0 \rightarrow G_n$ is the unit map.

Thanks to the above Lemma we have a cosimplicial diagram $(e_n^* \iota_{G_n}^* \psi_{G_n})_{n \geq 0}$ of morphisms of perfect modules on X , the limit of which is the morphism $q^* \iota_Y^* \psi_Y$. Thanks to the induction hypothesis every $e_n^* \iota_{G_n}^* \psi_{G_n}$ is an equivalence, and thus $q^* \iota_Y^* \psi_Y$ is an equivalence as well. One finally gets that $\iota_Y^* \psi_Y$ is an equivalence because q^* is conservative (q is a groupoid quotient morphism).

We get the result by induction on the degree of geometricity. \square

Remark 2.5. Just as in Remark 2.4 one can even prove that the equivalence ψ_X is homotopic to the identity.

Proof of the Lemma. The morphism ψ_X , resp. ψ_Y , is determined by the leading term of the isotropic fibration structure η_X , resp. η_Y . Hence the morphism $(\pi_Y^* f)^* \psi_Y$ is determined by the leading term of $(\pi_Y^* f)^* \eta_Y$, considered as an isotropic fibration structure on

$$f^* \pi_Y : f^* T^*[n]Y \longrightarrow X$$

and where we equip $f^* T^*[n]Y$ with the n -shifted presymplectic structure $(\pi_Y^* f)^* \omega_Y$. Now the result follows from the following fact:

- the pull-back of $(T^* f)^* \eta_X$ is naturally homotopic to $(\pi_Y^* f)^* \eta_Y$.

First of all this claim makes sense as we have already seen that the analogous statement for ω 's is true (see the proof of Lemma 2.3). It is even true for λ 's. Since the part of the above diagram we are interested in lives over X , all images of previously considered homotopies between 0 and images of λ 's in the space of 1-forms relative to X are themselves equivalent. The Lemma is proved. \square

End of the proof of Theorem 2.2. The first part of the Theorem follows from the third one. Namely, Lemma 1.3 tells us that if we have a non-degenerate isotropic fibration on an n -shifted presymplectic Artin stack then it is n -shifted symplectic. \square

2.3 A Lagrangian structure on shifted conormal stacks

Let $X \rightarrow Y$ be a morphism of Artin stacks. We have already seen that there is a tautological n -shifted 1-form on $T^*[n]Y$, mainly determined by the identity morphism of $T^*[n]Y$ (and the map $\pi_Y^* \mathbb{L}_Y[n] \rightarrow \mathbb{L}_{T^*[n]Y}$). We now want to consider the morphism of Artin stacks $\ell_f : T_X^*[n]Y \rightarrow T^*[n]Y$, where $T_X^*[n]Y := T^*[n-1]f = \mathbb{A}(\mathbb{L}_f[n-1])$ is the n -shifted conormal stack of f . First of all observe that this morphism factors through $f^* T^*[n]Y$, so that $\ell_f^* \omega_Y$ is homotopic to the pull-back of ω_X along the sequence

$$T_X^*[n]Y \rightarrow f^* T^*[n]Y \rightarrow T^*[n]X.$$

Hence $\ell_f^* \omega_Y$ is equivalent to 0 as the composed morphism of the above sequence factors through the zero section $X \rightarrow T^*[n]X$. Let us denote ω_f the isotropic structure we have just constructed on ℓ_f .

Our aim is to prove the following:

Theorem 2.8. *The n -shifted conormal stack is Lagrangian, that is to say the isotropic structure ω_f is non-degenerate.*

Let us provide two rather extreme examples:

- consider $f : X \rightarrow \text{pt}$, where pt is the terminal stack. In this case $\mathbf{T}_X^*[\mathbf{n}+1]\text{pt} \cong \mathbf{T}^*[\mathbf{n}]X$ has a Lagrangian morphism to the point equipped with its canonical $(\mathbf{n}+1)$ -shifted symplectic structure. We get back in a rather involved way the \mathbf{n} -shifted symplectic structure on an \mathbf{n} -shifted cotangent stack.
- consider the identity morphism $X \rightarrow X$. In this case $\mathbf{T}_X^*[\mathbf{n}]X \cong X$ and we get back that the zero section is Lagrangian.

2.3.1 Isotropic squares

Our aim is to apply a strategy similar to what we did in the previous Section. To do so we need to exhibit a structure on the projection $\mathbf{T}_X^*[\mathbf{n}]Y \rightarrow X$ that resembles to the one of a Lagrangian fibration. This is what we do in this Subsection.

Definition 2.9. An \mathbf{n} -shifted pre-isotropic square is commuting square of Artin stacks

$$\begin{array}{ccc} \mathbf{E} & \xrightarrow{g} & \mathbf{F} \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y \end{array}$$

together with the following additional structures:

- an \mathbf{n} -shifted presymplectic structure ω on \mathbf{F} : $\omega \in A^{2,\text{cl}}(\mathbf{F}, \mathbf{n})$.
- an isotropic structure γ on $\mathbf{E} \rightarrow \mathbf{F}$: $\gamma \in \text{Path}_{g^*\omega, 0}(A^{2,\text{cl}}(\mathbf{E}, \mathbf{n}))$.
- an isotropic fibration structure η on $\mathbf{F} \rightarrow Y$: $\eta \in \text{Path}_{\omega/Y, 0}(A^{2,\text{cl}}(\mathbf{F}/Y, \mathbf{n}))$.

Note that composing $\gamma_{/X}$ with $(g^*\eta)^{-1}$ one gets an element $\omega' \in \Omega_0 A^{2,\text{cl}}(\mathbf{E}/X, \mathbf{n}) \cong A^{2,\text{cl}}(\mathbf{E}/X, \mathbf{n}-1)$

Example 2.10. There is an \mathbf{n} -shifted pre-isotropic square structure with $\mathbf{F} = \mathbf{T}^*[\mathbf{n}]Y$, $\mathbf{E} = \mathbf{T}_X^*[\mathbf{n}]Y$, $g = \ell_f$, $\omega = \omega_Y$, $\gamma = \omega_f$ and $\eta = \eta_Y$.

Definition 2.11. Borrowing the above notation, an \mathbf{n} -shifted isotropic square is an \mathbf{n} -shifted pre-isotropic square equipped with a homotopy ξ in $A^{2,\text{cl}}(\mathbf{E}/X, \mathbf{n})$ between the two isotropic fibration structures $\gamma_{/X}$ and $g^*\eta$ on $\mathbf{E} \rightarrow X$.

We let the reader prove the following:

Proposition 2.12. The \mathbf{n} -shifted pre-isotropic square from Example 2.10 can be upgraded to an \mathbf{n} -shifted isotropic square. \square

2.3.2 Non-degeneracy of the isotropic structure on the \mathbf{n} -shifted conormal stack

Assume we are given an isotropic square. Borrowing the previous notation, we set $\mathbf{U} := \mathbf{T}_{\mathbf{E}}$, $\mathbf{V} := g^*\mathbf{T}_{\mathbf{F}/Y}$, $\mathbf{W} := g^*\mathbf{T}_{\mathbf{F}}$ and $\mathbf{Z} := \mathbf{T}_{\mathbf{E}/X}$. Recall that we have a lot of structure:

- a degree \mathbf{n} non-degenerate pairing on \mathbf{W} (given by ω).
- a path from 0 to its pull-back along $\mathbf{U} \rightarrow \mathbf{W}$ (given by γ).
- a non-degenerate path from 0 to its pull-back along $\mathbf{V} \rightarrow \mathbf{W}$ (given by η).
- these two paths compose to a self-homotopy of 0 in the space of degree \mathbf{n} pairings on $\mathbf{U} \times_{\mathbf{W}} \mathbf{V}$, defining a degree $\mathbf{n} - 1$ pairing β on it.
- a path from 0 to the pull-back of this last pairing along $\mathbf{Z} \rightarrow \mathbf{U} \times_{\mathbf{W}} \mathbf{V}$ (given by ξ).

Lemma 2.13. If ξ is non-degenerate, then so are β and γ

Proof. We have already seen that if ξ is non-degenerate then so is β (this is Lemma 1.3). Now the data of ω , γ and η provides us with the following commuting diagram

$$\begin{array}{ccccc}
\mathbf{U} \times_{\mathbf{W}} \mathbf{V} & \longrightarrow & \mathbf{V} & \longrightarrow & 0 \\
\downarrow & & \downarrow & & \downarrow \\
\mathbf{U} & \longrightarrow & \mathbf{W} & \longrightarrow & \mathbf{V}^\vee[\mathbf{n}] \\
\downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathbf{U}^\vee[\mathbf{n}] & \longrightarrow & (\mathbf{U} \times_{\mathbf{W}} \mathbf{V})^\vee[\mathbf{n}]
\end{array}$$

We observe that the following squares are Cartesian: top-left (by definition), top-right (by non-degeneracy of η), bottom-right (by definition), and the big one (by non-degeneracy of β). Therefore the bottom-left square is Cartesian as well, which means that γ is non-degenerate. \square

End of the proof of Theorem 2.8. Note that, if $\mathbf{E} = \mathbb{A}(\mathbf{E})$, $\mathbf{F} = \mathbb{A}(\mathbf{F})$ and g is linear in the fiber (i.e. g comes from a morphism $\mathbf{E} \rightarrow \mathbf{F}^*$) then we have that the cofiber of $\mathbf{Z} \rightarrow \mathbf{U} \times_{\mathbf{W}} \mathbf{V}$ is

$$\mathrm{cofib}(\pi_X^* \mathbf{E} \rightarrow \mathbb{T}_{\mathbf{E}} \times_{g^* \mathbb{T}_{\mathbf{F}}} g^* \mathbb{T}_{\mathbf{F}/\mathbf{Y}}) \cong \pi_X^* \mathbb{T}_{\mathbf{X}/\mathbf{Y}} = \pi_X^* \mathbb{T}_{\mathbf{f}}.$$

This is typically what happens in the situation of Example 2.10. In this case the null-homotopic sequence $\mathbf{Z} \rightarrow \mathbf{U} \times_{\mathbf{W}} \mathbf{V} \rightarrow \mathbf{Z}^\vee[\mathbf{n} - 1]$ then takes the following form:

$$\pi_X^* \mathbb{L}_{\mathbf{f}}[\mathbf{n} - 1] \longrightarrow \mathbf{U} \times_{\mathbf{W}} \mathbf{V} \longrightarrow \pi_X^* \mathbb{T}_{\mathbf{f}}.$$

Hence we get a map $\pi_X^* \mathbb{T}_{\mathbf{f}} \cong \mathrm{hocofib}(\mathbf{Z} \rightarrow \mathbf{U} \times_{\mathbf{W}} \mathbf{V}) \rightarrow \pi_X^* \mathbb{T}_{\mathbf{f}}$, which is an equivalence if and only if ξ is non-degenerate. As before, one can show that this map must be an equivalence by reasoning along the following steps:

1. using $\mathbb{G}_{\mathbf{m}}$ -equivariant enhancements, one proves (as in the proof of Lemma 2.5) that the map $\pi_X^* \mathbb{T}_{\mathbf{f}} \rightarrow \pi_X^* \mathbb{T}_{\mathbf{f}}$ is the pull-back along π_X^* of a map $\psi_{\mathbf{f}} : \mathbb{T}_{\mathbf{f}} \rightarrow \mathbb{T}_{\mathbf{f}}$.
2. one then proves (along the lines of Lemmas 2.3 and 2.6) that $\psi_{\mathbf{f}}$ is natural in the following sense: for any morphism commuting square

$$\begin{array}{ccc}
X' & \xrightarrow{f'} & Y' \\
x \downarrow & & \downarrow y \\
X & \xrightarrow{f} & Y
\end{array}$$

one has a commuting square

$$\begin{array}{ccc}
\mathbb{T}_{f'} & \xrightarrow{\psi_{f'}} & \mathbb{T}_{f'} \\
\downarrow & & \downarrow \\
x^* \mathbb{T}_f & \xrightarrow{x^* \psi_f} & x^* \mathbb{T}_f
\end{array}$$

3. the previous step allows one to reduce to the case when both X and Y are affine, by induction on the degree of geometricity of f (as in Corollaries 2.4 and 2.7).

Remark 2.6. We say that f is \mathbf{m} -geometric if it can be realized as the colimit $f = |f_*|$ of a simplicial diagram $f_* : X_* \rightarrow Y_*$ of morphisms, with both X_* and Y_* being Segal groupoids in $(\mathbf{m} - 1)$ -geometric stacks. Every morphism between geometric stacks is \mathbf{m} -geometric for some \mathbf{m} . Namely, Y being geometric there exists a Segal groupoid Y_* such that $Y = |Y_*|$ and every Y_n is geometric. Then there exists a Segal groupoid X_* presenting X such that $X_n \cong X \times_Y Y_n$ and a simplicial morphism f_* presenting f such that f_n is equivalent to the second projection (see [HAG2, §1.3.5]). Observe that every X_n is geometric, as geometricity is preserved by fiber products.

4. one finally proves that the result is true when both X and Y are affine.

We conclude by using Lemma 2.13: since ξ is non-degenerate then so is $\gamma = \omega_f$. \square

Remark 2.7. We won't do it but one can also prove, using similar methods, that given two morphisms $X_1 \rightarrow Y$ and $X_2 \rightarrow Y$, the morphism

$$X_1 \times_Y X_2 \rightarrow \mathbf{T}_{X_1}^*[n]Y \times_{\mathbf{T}^*[n]Y} \mathbf{T}_{X_2}^*[n]Y$$

has a Lagrangian structure.

3 Shifted cotangent stacks in derived Poisson geometry

In this Section we provide several conjectural relations between shifted cotangent stacks and shifted Poisson structures ([CPT+, Pr]).

3.1 The n -shifted Poisson structure of $\mathbf{T}^*[n]X$ and shifted polyvector fields

On the one hand we know from [CPT+, Pr] that the spaces of n -shifted symplectic structures and of non-degenerate n -shifted Poisson structures (we refer to *loc. cit.* for a precise definition, and follow the notation from [CPT+]) on a given Artin stack are equivalent. We therefore obtain an n -shifted Poisson structure on $\mathbf{T}^*[n]X$, and thus in particular a \mathbb{P}_{n+1} -algebra structure extending the natural unbounded cda structure of

$$\Gamma(\mathbf{T}^*[n]X, \mathcal{O}_{\mathbf{T}^*[n]X}) = \Gamma(X, \mathrm{Sym}(\mathbb{T}_X[-n])).$$

Moreover, recall that $\mathbf{T}^*[n]X$ carries a \mathbb{G}_m -action, which makes $\Gamma(X, \mathrm{Sym}(\mathbb{T}_X[-n]))$ a commutative algebra object in graded complexes (the weight being obviously given by the symmetric degree). We have seen that the n -shifted symplectic structure on $\mathbf{T}^*[n]X$ is homogeneous of weight 1 for this \mathbb{G}_m -action.

One can carry over the \mathbb{G}_m -action along [CPT+] (this ends up working relatively to $B\mathbb{G}_m$, and the constructions from *loc. cit.* do extend to the relative setting) and get a $\mathbb{P}_{n+1}^{\mathrm{gr}}$ -algebra structure on $\Gamma(X, \mathrm{Sym}(\mathbb{T}_X[-n]))$. Recall that $\mathbb{P}_{n+1}^{\mathrm{gr}}$ for a graded version of \mathbb{P}_{n+1} where the Poisson bracket generator has weight -1 .

We believe the following is true:

Conjecture 3.1. *The above $\mathbb{P}_{n+1}^{\mathrm{gr}}$ -algebra structure on $\Gamma(X, \mathrm{Sym}(\mathbb{T}_X[-n]))$ is equivalent to the $\mathbb{P}_{n+1}^{\mathrm{gr}}$ -algebra $\mathrm{Pol}(X, n)$ of n -shifted polyvector fields from [CPT+].*

3.2 Lagrangian morphisms, deformed shifted cotangent stacks and shifted Poisson structures

Let $f : X \rightarrow Y$ be a morphism of Artin stacks equipped with an n -shifted Lagrangian structure γ (in particular, Y implicitly carries an n -shifted symplectic structure ω). Note that the leading term γ_0 of γ provides us with an equivalence $f^*\mathbb{T}_f[1] \rightarrow \mathbb{L}_X[n]$, which leads to an equivalence of derived stacks $\mathbf{T}_X Y := \mathbb{A}(f^*\mathbb{T}_f[1]) \rightarrow \mathbf{T}^*[n]$.

We then consider the formal completion $\widehat{Y}_f = X_{\mathrm{DR}} \times_{Y_{\mathrm{DR}}} Y$ of Y along f . The morphism f factors as $X \xrightarrow{\widehat{f}} \widehat{Y}_f \rightarrow Y$, and one can easily see that

- the pull-back $\widehat{\omega}$ of ω along $\widehat{Y}_f \rightarrow Y$ is still non-degenerate, thus defining an n -shifted symplectic structure on \widehat{Y}_f .
- the homotopy γ between $f^*\omega = \widehat{f}^*\widehat{\omega}$ and 0 , seen as an isotropic structure on \widehat{f} , is still non-degenerate, thus defining an n -shifted Lagrangian structure on \widehat{f} .

As a matter of notation, if E is a perfect module over X then we simply write \widehat{E} for the formal completion of E along the zero section. It follows from [GR, Chapter IV.5] that there actually exists a morphism $\widehat{f} : X \times [\mathbb{A}^1/\mathbb{G}_m] \rightarrow \widehat{Y}_f$ of derived stacks over $[\mathbb{A}^1/\mathbb{G}_m]$ such that

- the fiber at 0 is the zero section $X \rightarrow \widehat{T}_X Y \cong \widehat{T}^*[n]X$.
- the fiber at 1 is $\widehat{f}: X \rightarrow \widehat{Y}_f$.

As a consequence we get that there is a filtration on the unbounded cdga $\Gamma(\widehat{Y}_f, \mathcal{O}_{\widehat{Y}_f})$ with associated graded being $\Gamma(\widehat{T}^*[n]X, \mathcal{O}_{\widehat{T}^*[n]X}) = \Gamma(X, \text{Sym}(\mathbb{T}_X[-n]))$. In other words, we get that $\Gamma(X, \text{Sym}(\mathbb{T}_X[-n]))$ becomes equipped with a mixte differential.

Remark 3.1. Note that there is a much easier way to see this. Thanks to the equivalence $f^*\mathbb{T}_f[1] \rightarrow \mathbb{L}_X[n]$ provided by γ_0 , we get an equivalence

$$\Gamma(X, \text{Sym}(\mathbb{T}_X[-n])) \rightarrow \Gamma(X, \text{Sym}(f^*\mathbb{L}_f[-1])) \cong \mathbf{DR}(X/Y)^\sharp$$

of commutative algebra objects in graded complexes. The mixte differential we get on $\Gamma(X, \text{Sym}(\mathbb{T}_X[-n]))$ then comes from the mixte differential on $\mathbf{DR}(X/Y)$.

We believe the following is true:

Conjecture 3.2. a) *There exists a relative n -shifted symplectic structure $\widetilde{\omega}$ on \widehat{Y}_f over $[\mathbb{A}^1/\mathbb{G}_m]$ such that $1^*\widetilde{\omega} \cong \widehat{\omega}$ and $0^*\widetilde{\omega} \cong \omega_X$.*
b) *There exists a relative Lagrangian structure $\widetilde{\gamma}$ on \widehat{f} (over $[\mathbb{A}^1/\mathbb{G}_m]$) such that $1^*\widetilde{\gamma} \cong \gamma$ and $0^*\widetilde{\gamma} \cong \gamma_X$.*

Assuming the above conjecture and working out the constructions from [CPT+] relatively to $[\mathbb{A}^1/\mathbb{G}_m]$ we get that the mixte differential on $\Gamma(X, \text{Sym}(\mathbb{T}_X[-n]))$ is compatible with the \mathbb{P}_{n+1}^{gr} -algebra. We believe that the following is also true:

Conjecture 3.3. *The above mixte differential is given by the bracket with a weight 2 Maurer Cartan element π_f in the graded dg Lie algebra $\Gamma(X, \text{Sym}(\mathbb{T}_X[-n]))[n]$.*

Putting together Conjectures 3.1, 3.2 and 3.3 we obtain a map from n -shifted Lagrangian morphisms $f: X \rightarrow Y$ to $(n-1)$ -shifted Poisson structures π on X . We don't expect this map $f \mapsto \pi_f$ to be an equivalence as the construction only depends on \widehat{f} (namely, $\pi_f = \pi_{\widehat{f}}$). But we expect the following

Conjecture 3.4. *The map $f \mapsto \pi_f$ provides an equivalence between:*

- *the space of n -shifted Lagrangian morphisms $f: X \rightarrow Y$ such that $\widehat{Y}_f \rightarrow Y$ is an equivalence.*
- *the space of $(n-1)$ -shifted Poisson structures on X in the sense of [CPT+].*

References

- [Ca] D. Calaque, Lagrangian structures on mapping stacks and semi-classical TFTs, in *Stacks and Categories in Geometry, Topology, and Algebra* (Pantev et al. editors), Contemporary Mathematics **643** (2015).
- [CPT+] D. Calaque, T. Pantev, B. Toën, M. Vaquié & G. Vezzosi, Shifted Poisson Structures and Deformation Quantization, preprint [arXiv:1506.03699](https://arxiv.org/abs/1506.03699).
- [GR] D. Gaitsgory & N. Rozenblyum, *A study in derived algebraic geometry*, book in progress, preliminary version available at <http://www.math.harvard.edu/~gaitsgde/GL/>
- [HAG2] B. Toën & G. Vezzosi, Homotopical Algebraic Geometry II: geometric stacks and applications, *Memoirs of the American Mathematical Society* 2008, 224 pp.
- [Ha] R. Haugseng, Iterated spans and “classical” topological field theories, preprint [arXiv:1409.0837](https://arxiv.org/abs/1409.0837).
- [Me] V. Melani, Poisson bivectors and Poisson brackets on affine derived stacks, *Adv. in Math.* **288** (2016), 1097–1120.

- [PTVV] T. Pantev, B. Toën, M. Vaquié & G. Vezzosi, Shifted Symplectic Structures, Publications mathématiques de l’IHÉS **117** (2013), no. 1, 271–328.
- [Pr] J.P. Pridham, Shifted Poisson and symplectic structures on derived N-stacks, preprint [arXiv:1504.01940](https://arxiv.org/abs/1504.01940).
- [Sa] P. Safronov, Quasi-Hamiltonian reduction via classical Chern–Simons theory, Adv. in Math. **287** (2016), 733–773.